

Contec® Professional Guide to Proper Wiping

How to Saturate and Clean with ContecCloth™ Cloth BLUE Wipes:



1. Assemble the bucket (PRCN0005) by attaching the lid.
2. Remove the tab around the lid.
3. Label bucket with the name of the appropriate facility approved disinfectant.



1. Count out a stack of approximately 25 wipes (PRMW1213A).
2. Place the stack of wipes into the bucket.



1. Fill a 32 oz bottle with properly diluted facility approved disinfectant
2. Carefully pour the bottle of disinfectant over stack of wipes; be sure to saturate all corners.
3. Allow about 1-2 minutes for the wipes to absorb the disinfectant and become fully saturated.



1. Wipe surfaces with unidirectional strokes, going in the same direction each time.
2. Slightly overlap each stroke to ensure even coverage of disinfectant and avoid creating any missed areas.



1. Use all clean sides of the wipe. The natural quarterfold creates 8 clean, fresh sides of each wipe.
2. Get a new wipe when the wipe no longer dispenses even fluid. Always get new wipe becomes excessively soiled.
3. Always use different wipes for patient bathroom.



1. Dispose the used, soiled wipe in the trash.
2. Always follow the proper biohazard waste disposal procedures (blood, bodily fluid, etc.)

DISCLAIMER: The following document is intended to serve as a tool to provide documentation of competency only and is in no way providing certification of the information in this document. Facilities should follow their own guidelines, policies and procedures which, in all cases, will take precedence over this training document.